

Technical Data Sheet



Retatrutide

Cat No.:	HY-P3506
CAS No.:	2381089-83-2
Molecular Formula:	C ₂₂₁ H ₃₄₂ N ₄₆ O ₆₈
Molecular Weight:	4731.33
Sequence:	Tyr-{Aib}-Gln-Gly-Thr-Phe-Thr-Ser-Asp-Tyr-Ser-Ile- α -Me-Leu-Leu-Asp-Lys-(diacid-C2 0-gamma-Glu-(AEEA)-Lys)-Ala-Gln-{Aib}-Ala-Phe-Ile-Glu-Tyr-Leu-Leu-Glu-Gly-Gly-Pro- Ser-Ser-Gly-Ala-Pro-Pro-Pro-Ser-NH ₂
Sequence Shortening:	Y-{Aib}-QGTFTSDYSI- α -Me-Leu-LDK-{Lys(AEEA- γ -Glu-C20 diacid)}AQ-{Aib}-AFIEYLLEG GPSSGAPPPS-NH ₂
Target:	GCGR; GLP Receptor
Pathway:	GPCR/G Protein
Storage:	Sealed storage, away from moisture and light, under nitrogen Powder -80°C 2 years -20°C 1 year * In solvent : -80°C, 6 months; -20°C, 1 month (sealed storage, away from moisture and light, under nitrogen)

Tyr-{Aib}-Gln-Gly-Thr-Phe-Thr-Ser-Asp-
Tyr-Ser-Ile- α -Me-Leu-Leu-Asp-Lys-
{diacid-C20- γ -Glu-(AEEA)-Lys)-Ala-Gln-
{Aib}-Ala-Phe-Ile-Glu-Tyr-Leu-Leu-Glu-
Gly-Gly-Pro-Ser-Ser-Gly-Ala-Pro-Pro-
Pro-Ser-NH₂

SOLVENT & SOLUBILITY

In Vitro	DMSO : 50 mg/mL (10.57 mM; Need ultrasonic) H ₂ O : 20 mg/mL (4.23 mM; ultrasonic and adjust pH to 9 with NH ₃ -H ₂ O)
----------	--

	Solvent Mass Concentration	1 mg	5 mg	10 mg
Preparing Stock Solutions	1 mM	0.2114 mL	1.0568 mL	2.1136 mL
	5 mM	0.0423 mL	0.2114 mL	0.4227 mL
	10 mM	0.0211 mL	0.1057 mL	0.2114 mL

Please refer to the solubility information to select the appropriate solvent.

BIOLOGICAL ACTIVITY

Description	Retatrutide (LY3437943) is a triple agonist peptide of the glucagon receptor (GCGR), glucosdependent insulinotropic polypeptide receptor (GIPR), and glucagon-like peptide-1 receptor (GLP-1R). Retatrutide binds human GCGR, GIPR, and GLP-1R with EC ₅₀ values of 5.79, 0.0643 and 0.775 nM, respectively. Retatrutide can be used for the research of obesity[1].
IC ₅₀ & Target	EC ₅₀ (for human): 5.79 (GCGR), 0.0643 (GIPR), 0.775 nM (GLP-1R)[1]. EC ₅₀ (for mouse): 2.32 (GCGR), 0.191 (GIPR), 0.794 nM (GLP-1R)[1]. Ki (for human): 5.6 (GCGR), 0.057 (GIPR), 7.2 nM (GLP-1R)[1]. Ki (for mouse): 73 (GCGR), 2.8 (GIPR), 1.3 nM (GLP-1R)[1].

In Vitro	Retatrutide (LY3437943) has efficacy for human GCGR, GIPR, and GLP-1R with EC ₅₀ values of 5.79, 0.0643 and 0.775 nM, respectively[1].						
	Retatrutide has efficacy for mouse GCGR, GIPR, and GLP-1R with EC ₅₀ values of 2.32, 0.191 and 0.794 nM, respectively[1].						
	Retatrutide has binding affinity for human GCGR, GIPR, and GLP-1R with K _i values of 5.6, 0.057 and 7.2 nM, respectively[1].						
	Retatrutide has binding affinity for mouse GCGR, GIPR, and GLP-1R with K _i values of 73, 2.8 and 1.3 nM, respectively[1].						
In Vivo	Retatrutide (LY3437943) (s.c.; 47 µg/kg; single) engages GCGR in vivo and can improve glucose tolerance in an ipGTT through either the GIP or GLP-1 receptors[1].						
	Retatrutide (s.c.; 10 mL/kg; cycle every 3 days; for 21 days) causes great body weight loss and increases energy expenditure through glucagon receptor activation[1].						
	Retatrutide has safety and tolerability[1].						
	Animal Model:	Male CD-1 mice[1]					
	Dosage:	47 µg/kg					
	Administration:	Subcutaneous administration, single					
	Result:						
		AUC _{last} , ng*h/mL	AUC _{0-∞} , ng*h/mL	C _{max} , ng/mL	T _{max} , h	t _{1/2} , h	CLF, mL/h/kg
		41135	41905	1680	12	21	11.22
	Animal Model:	Diet-induced obese (DIO) male C57/Bl6 mice (24-25 weeks, 40-51 g)[1]					
	Dosage:	10 mL/kg					
	Administration:	Subcutaneous (SC) injection, cycle every 3 days, for 21 days					
	Result:	Decreased body weight and improved glycemic control.					